

References related to Table S3 from Schorderet et al (KPSB #1471299)

44. Kardailsky I, Shukla VK, Ahn JH, Dagenais N, Christensen SK, Nguyen JT, Chory J, Harrison MJ, Weigel D. Activation tagging of the floral inducer FT. *Science*. 1999;286:1962–1965.
45. Mandel MA, Gustafson-Brown C, Savidge B, Yanofsky MF. Molecular characterization of the Arabidopsis floral homeotic gene APETALA1. *Nature*. 1992;360:273–277. doi:10.1038/360273a0.
46. Bradley D, Ratcliffe O, Vincent C, Carpenter R, Coen E. Inflorescence commitment and architecture in Arabidopsis. *Science*. 1997;275:80–83.
47. Hartmann U, Hohmann S, Nettekheim K, Wisman E, Saedler H, Huijser P. Molecular cloning of SVP: a negative regulator of the floral transition in Arabidopsis. *Plant J*. 2000;21:351–360.
48. Golz JF, Keck EJ, Hudson A. Spontaneous mutations in KNOX genes give rise to a novel floral structure in Antirrhinum. *Curr Biol*. 2002;12:515–522.
49. Kieffer M, Stern Y, Cook H, Clerici E, Maulbetsch C, Laux T, Davies B. Analysis of the transcription factor WUSCHEL and its functional homologue in Antirrhinum reveals a potential mechanism for their roles in meristem maintenance. *Plant Cell*. 2006;18:560–573.
50. Huijser P, Klein J, Lonnig WE, Meijer H, Saedler H, Sommer H. Bracteomania, an inflorescence anomaly, is caused by the loss of function of the MADS-box gene SQUAMOSA in Antirrhinum majus. *EMBO J*. 1992;11:1239–1249.
51. Bradley D, Carpenter R, Copsey L, Vincent C, Rothstein S, Coen E. Control of inflorescence architecture in Antirrhinum. *Nature*. 1996;379:791–797.
52. Masiero S, Li MA, Will I, Hartmann U, Saedler H, Huijser P, Scharz-Sommer Z, Sommer H. INCOMPOSITA: a MADS-box gene controlling prophyll development and floral meristem identity in Antirrhinum. *Development*. 2004;131:5981–5990.
53. Stuurman J, Jäggi F, Kuhlmeier C. Shoot meristem maintenance is controlled by a GRAS-gene mediated signal from differentiating cells. *Genes Dev*. 2002;16:2213–2218.
54. Tsukamoto A, Hirai T, Chin DP, Mii M, Mizoguchi T, Mizuta D, Yoshida H, Olsen JE, Ezura H, Fukuda N. The FT-like gene PehFT in petunia responds to photoperiod and light quality but is not the main gene promoting light quality associated flowering. *Plant Biotechnol J*. 2016;33:297–307.
55. Molinero-Rosales N, Jamilena M, Zurita S, Gomez P, Capel J, Lozano R. *FALSIFLORA*, the tomato orthologue of *FLORICAULA* and *LEAFY*, controls flowering time and floral meristem identity. *Plant J*. 1999;20:685–693.
56. Janssen B-J, Williams A, Chen J-J, Mathern J, Hake S, Sinha N. Isolation and characterization of two knotted-like homeobox genes from tomato. *Plant Mol. Biol*. 1998;36:417–425.
57. Reinhardt D, Frenz M, Mandel T, Kuhlmeier C. Microsurgical and laser ablation analysis of interactions between the zones and layers of the tomato shoot apical meristem. *Development*. 2003;130:4073–4083.
58. Krieger U, Lippman ZB, Zamir D. The flowering gene SINGLE FLOWER TRUSS drives heterosis for yield in tomato. *Nat Genet*. 2010;42:459–U138.
59. Burko Y, Shleizer-Burko S, Yanai O, Schwartz I, Zelnik ID, Jacob-Hirsch J, Kela I, Eshed-Williams L, Ori N. A role for APETALA1/FRUITFULL transcription factors in tomato leaf development. *Plant Cell*. 2013;25:2070–2083.
60. Pnueli L, Carmel-Goren L, Hareven D, Gutfinger T, Alvarez J, Ganai M, Zamir D, Lifschitz E. The SELF-PRUNING gene of tomato regulates vegetative to reproductive switching of sympodial meristems and is the ortholog of CEN and TFL1. *Development*. 1998;125:1979–1989.
61. Nakano T, Kato H, Shima Y, Ito Y. Apple SVP family MADS-Box proteins and the tomato pedicel abscission zone regulator JOINTLESS have similar molecular activities. *Plant Cell Physiol*. 2015;56:1097–1106.
62. Hofer J, Turner L, Hellens R, Ambrose M, Matthews P, Michael A, Ellis N. UNIFOLIATA regulates leaf and flower morphogenesis in pea. *Curr Biol*. 1997;7:581–587.
63. Taylor S, Hofer J, Murfet I. STAMINA PISTILLOIDA, the pea ortholog of FIM and UFO, is required for normal development of flowers, inflorescences, and leaves. *Plant Cell*. 2001;13:31–46.
64. Hofer J, Gourlay C, Michael A, Ellis THN. Expression of a class 1 knotted1-like homeobox gene is down-regulated in pea compound leaf primordia. *Plant Mol Biol*. 2001;45:387–398.
65. Hecht V, Laurie RE, Vander Schoor JK, Ridge S, Knowles CL, Liew LC, Sussmilch FC, Murfet IC, Macknight RC, Weller JL. The pea *GIGAS* gene is a *FLOWERING LOCUS T* homolog necessary for graft-transmissible specification of flowering but not for responsiveness to photoperiod. *Plant Cell*. 2011;23:147–161.
66. Taylor SA, Hofer JMI, Murfet IC, Sollinger JD, Singer SR, Knox MR, Ellis THN. PROLIFERATING INFLORESCENCE MERISTEM, a MADS-box gene that regulates floral meristem identity in pea. *Plant Physiol*. 2002;129:1150–1159.

67. Foucher F, Morin J, Courtiade J, Cadioux S, Ellis N, Banfield MJ, Rameau C. *DETERMINATE* and *LATE FLOWERING* are two *TERMINAL FLOWER1/CENTRORADIALIS* homologs that control two distinct phases of flowering initiation and development in pea. *Plant Cell*. 2003;15:2742-2754.